

Drinking and Metabolic Disorders: Type 2 Diabetes Mellitus and Metabolic Syndrome

BACKGROUND

The liver and the pancreas are responsible for producing the enzymes involved in breaking down the carbohydrates, fats, and proteins in food and turning them into energy. If these metabolites are not used immediately, the body stores them in the muscles, fat cells, or liver for later use.

Metabolic disorders is an umbrella term for conditions in which food metabolism is disrupted, either through problems in the liver or pancreas, or in the utilization of stored energy.

This *IARD Health Review* focuses on the two most common metabolic disorders, Type 2 diabetes mellitus and MetS.

The International Diabetes Federation estimates the prevalence of diabetes worldwide to be 8.3% [2], with Type 2 diabetes accounting for 90% of all cases [3].

- The global burden of disease attributable to diabetes in 2012 was 1.5 million deaths and 89 million disability adjusted life years (DALYs) [4].
- Individuals with Type 2 diabetes are at increased risk for kidney failure, stroke, cardiovascular disease, diabetic retinopathy, and amputation due to nerve damage and poor circulation to the feet [5].

MetS is estimated to affect at least 20% to 30% of the adult population in most countries [6]. Rates are higher in Latin America than in the U.S. and Canada or across Europe.

- The global burden of disease attributable to MetS is difficult to assess, as each of the defining components of MetS is a risk factor for cardiovascular disease, and individuals with MetS often develop diabetes [7].
- Individuals with MetS are at increased risk for atherosclerotic cardiovascular disease and Type 2 diabetes [8].

It is widely acknowledged that the increasing rate of obesity worldwide is driving an increasing rate in MetS, which in turn contributes to an increasing rate of Type 2 diabetes [9, 10] and cardiovascular disease [11].

Diabetes risk is also linked to genetic factors, age, ethnicity, and in some populations maternal malnutrition.

SUMMARY OF THE EVIDENCE

The relationship between alcohol intake and risk for Type 2 diabetes is characterized by a J-shaped or a U-shaped curve, similar to what is observed for cardiovascular disease and all-cause mortality.

- Moderate drinkers are at lower risk for Type 2 diabetes than abstainers and heavier drinkers [12-14].
- This relationship has been observed in both men and women [15, 16] and holds true across cultures and ethnic groups [17-19].
- Heavy drinking and heavy episodic drinking are associated with increased risk for Type 2 diabetes [20-25].

The relationship between MetS and alcohol consumption also follows a J-shaped curve.

- A protective effect has been described for moderate drinking and risk of MetS [22, 26-29], with some evidence suggesting that the benefit may be related to drinking in conjunction with meals [30, 31].
- Heavy drinking is associated with an increase in risk for MetS [19, 22, 32], as is heavy episodic drinking, particularly among occasional heavy drinkers [33].

MetS and Type 2 diabetes contribute to risk for other serious or life-threatening conditions and diseases.

- Individuals with Type 2 diabetes are at increased risk for kidney failure [4], stroke and cardiovascular disease (CVD) [34], diabetic retinopathy, and amputation.
- Individuals with MetS may have twice the risk for cardiovascular disease [11, 35] and five times the risk for Type 2 diabetes [36] compared with individuals without the syndrome.
- The cardioprotective benefits of moderate alcohol consumption may be especially important in metabolic disorder patients because of the prevalence of cardiovascular comorbidities within these populations [27].
- As alcohol intake increases, cardiovascular protection may be offset by increased risk for hypertension [37].

Mechanisms of action

The biological mechanisms underlying the relationship between moderate drinking and Type 2 diabetes and MetS have been well described.

- Alcohol consumption appears to improve insulin sensitivity without affecting insulin secretion [16, 38, 39], which helps to regulate plasma glucose levels [40]. High plasma glucose levels are a diagnostic marker for both Type 2 diabetes and MetS.
- Light-to-moderate alcohol consumption is associated with more favorable measures of fasting triglycerides, fasting glucose, and hemoglobin A1c among obese subjects [41] and among broader populations that include subjects with and without MetS [42].
- Moderate alcohol consumption is associated with an elevated blood concentration of high-density lipoprotein (HDL) [43-45]. Low HDL levels are associated with MetS and cardiovascular disease risk [27].

Among individuals with Type 2 diabetes or MetS, moderate consumption of alcohol may reduce the risk for other related diseases and improve disease outcomes.

- Moderate drinking appears to reduce the risk of cardiovascular disease [46] and cardiovascular deaths [14], and deaths from all causes among Type 2 diabetes patients [46].
- Elevated HDL cholesterol concentration, lower insulin levels and a lower risk of ischemic heart disease have been reported in association with moderate drinking among men, both with and without MetS [27].
- A decrease in the progression of atherosclerosis has been demonstrated in moderate drinkers with Type 2 diabetes [47].
- Heavy drinking, conversely, increases blood pressure [48] and triglycerides [49], both of which are among the features of MetS.

Daily moderate drinking as part of a high quality diet may improve Type 2 diabetes and MetS outcomes.

- A Mediterranean-type diet that includes daily moderate drinking has been associated with reduced cardiovascular mortality and all-cause mortality among patients with Type 2 diabetes. Furthermore, daily moderate drinking is reported to have the largest preventive impact on overall mortality among other identified components of the Mediterranean diet [50].
- Similarly, among patients with MetS, adherence to a Mediterranean-type diet that includes daily moderate drinking was associated with lower prevalence of MetS, progression of MetS, and several measures of individual MetS components, including abdominal adiposity, HDL cholesterol, triglycerides, fasting glucose, and insulin resistance [42].

There is evidence that harmful drinking patterns may interfere with treatment compliance among Type 2 diabetes patients [23] and worsen Type 2 diabetes outcomes [51].

Influences on the relationship between alcohol consumption and metabolic disorders

The relationship between drinking and metabolic disorders is moderated by age, gender, body mass index (BMI), diet quality, and physical activity.

- Age plays a role in risk of onset of Type 2 diabetes and MetS, as insulin resistance and glucose intolerance increase with age.
- However, the relationship between Type 2 diabetes and moderate drinking seems particularly robust in older adults [52].
- Some studies find that the protective effects of moderate alcohol consumption are greater for women than for men [24], with more significant reductions in risk for Type 2 diabetes.
- Obesity plays a substantial role in the development and progression of Type 2 diabetes [17, 53], and is a defining criterion for MetS.
- Physical activity significantly reduces the risk for obesity, MetS, and Type 2 diabetes. Among patients with MetS, moderate drinking combined with physical activity has a greater effect on MetS than moderate drinking alone [42].

METHODOLOGICAL CONSIDERATIONS

When interpreting research findings on risk factors and associated health outcomes, a number of methodological issues should be taken into consideration.

Confounding:

Confounding variables can obscure the true nature of the relationship between drinking and metabolic disorders, and need to be identified and controlled to the greatest extent possible.

- Age, gender, ethnicity, and weight/obesity status, as indicated by BMI, are differentially associated with both drinking and with metabolic disorders, or, more specifically, with conditions strongly related to a metabolic disorder, and may confound the relationship with drinking.

Underreporting:

Underreporting of consumption is a central concern of alcohol-related research. It is well established that survey respondents often underestimate their alcohol consumption. The magnitude of underreporting varies by respondent, context, and the approach used to measure consumption.

Misclassification of Drinkers:

Studies suggest that the underreporting of consumption can result in the misclassification of drinkers, especially between the low and moderate consumption categories. This misclassification can make it difficult to establish a clear relationship between low or moderate alcohol intake and disease outcomes. This, in turn, makes it difficult to identify definitive thresholds at which alcohol consumption increases the risk for specific diseases.

TERMS AND CONCEPTS

Insulin is a hormone, produced by the pancreas, which is released in response to an increase in blood glucose levels. Insulin controls the transport of glucose through the bloodstream into and out of the muscles, fat cells and other tissues in response to high or low glucose levels in the blood.

Diabetes mellitus refers to a class of diseases in which blood sugar levels are elevated over long periods of time, either through the failure of the pancreas to produce sufficient insulin or of cells to respond to insulin stimulation. There are three types of diabetes:

- Type 1 diabetes mellitus, formerly called insulin dependent diabetes, occurs when the pancreas loses its ability to produce sufficient insulin, and usually manifests itself in childhood or young adulthood.
- Type 2 diabetes mellitus, formerly called non-insulin dependent diabetes, occurs when tissue or red blood cells become resistant to insulin, often accompanied by reduced insulin secretion by the pancreas. Type 2 diabetes is also referred to as adult-onset diabetes, usually manifesting itself in adulthood.
- Gestational diabetes is a temporary condition of insulin resistance occurring during pregnancy to women with no previous history of diabetes. Women who have had Gestational diabetes have a higher risk of developing Type 2 diabetes later in life.

Metabolic syndrome (MetS) is a widespread disorder of the metabolic processes governing energy utilization and storage. Diagnosis requires three out of the following five correlated conditions: abdominal obesity, high blood pressure, high fasting plasma glucose levels, high serum triglycerides, and low levels of high-density (HDL) cholesterol.

Ranges for moderate and heavy drinking and metabolic disorders:

There is little consistency in the definitions used across research studies regarding “moderate” and “heavy” alcohol consumption. In some instances, definitions may include average volume, average frequency of alcohol consumption, or a combination of both. This lack of consensus on what constitutes moderate or heavy drinking can diminish cross-study comparability.

REFERENCES

1. United Nations General Assembly. (2014). *Report of the open working group of the general assembly on sustainable development goals* (No. document A/68/970).
2. International Diabetes Foundation. (2014). *IDF Diabetes Atlas*. Retrieved from https://www.idf.org/sites/default/files/Atlas-poster-2014_EN.pdf
3. World Health Organization. (1999). *Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: Diagnosis and classification of diabetes mellitus*. Retrieved
4. World Health Organization (WHO). (2011). *Global status report on noncommunicable diseases 2010*. Geneva: World Health Organization (WHO).
5. World Health Organization (WHO). (2015). *Diabetes Fact Sheet*. Retrieved from <http://www.who.int/mediacentre/factsheets/fs312/en/>
6. Grundy, S. M. (2008). Metabolic syndrome pandemic. *Arterioscler Thromb Vasc Biol*, 28(4), 629-636.
7. International Diabetes Federation. (2006). *The IDF consensus worldwide definition of Metabolic Syndrome*. Retrieved from http://www.idf.org/webdata/docs/IDF_Meta_def_final.pdf
8. American Heart Association. (2012). *What is Metabolic Syndrome?* Retrieved from http://www.heart.org/idc/groups/heart-public/@wcm/@hcm/documents/downloadable/ucm_300322.pdf
9. Bergman, R. N., Van Citters, G. W., Mittelman, S. D., Dea, M. K., Hamilton-Wessler, M., Kim, S. P., et al. (2001). Central role of the adipocyte in the metabolic syndrome. *Journal of Investigative Medicine*, 49(1), 119-126.
10. Grundy, S. M. (2004). Obesity, metabolic syndrome, and cardiovascular disease. *Journal of Clinical Endocrinology and Metabolism*, 89(6), 2595-2600.
11. Isomaa, B., Almgren, P., Tuomi, T., Forsen, B., Lahti, K., Nissen, M., et al. (2001). Cardiovascular morbidity and mortality associated with the metabolic syndrome. *Diabetes Care*, 24(4), 683-689.
12. Koppes, L. L., Dekker, J. M., Hendriks, H. F., Bouter, L. M., & Heine, R. J. (2005). Moderate alcohol consumption lowers the risk of type 2 diabetes: a meta-analysis of prospective observational studies. *Diabetes Care*, 28(3), 719-725.
13. Baliunas, D. O., Taylor, B. J., Irving, H., Roerecke, M., Patra, J., Mohapatra, S., et al. (2009). Alcohol as a risk factor for type 2 diabetes: A systematic review and meta-analysis. *Diabetes Care*, 32(11), 2123-2132.
14. Howard, A. A., Arnsten, J. H., & Gourevitch, M. N. (2004). Effect of alcohol consumption on diabetes mellitus: a systematic review. *Ann Intern Med*, 140(3), 211-219.
15. Beulens, J. W., Stolk, R. P., van der Schouw, Y. T., Grobbee, D. E., Hendriks, H. F., & Bots, M. L. (2005). Alcohol consumption and risk of type 2 diabetes among older women. *Diabetes Care*, 28(12), 2933-2938.
16. Englund Ogge, L., Brohall, G., Behre, C. J., Schmidt, C., & Fagerberg, B. (2006). Alcohol consumption in relation to metabolic regulation, inflammation, and adiponectin in 64-year-old Caucasian women: a population-based study with a focus on impaired glucose regulation. *Diabetes Care*, 29(4), 908-913.
17. Lapidus, L., Bengtsson, C., Bergfors, E., Bjorkelund, C., Spak, F., & Lissner, L. (2005). Alcohol intake among women and its relationship to diabetes incidence and all-cause mortality: the 32-year follow-up of a population study of women in Gothenburg, Sweden. *Diabetes Care*, 28(9), 2230-2235.
18. Nakanishi, N., Suzuki, K., & Tataru, K. (2003). Alcohol consumption and risk for development of impaired fasting glucose or type 2 diabetes in middle-aged Japanese men. *Diabetes Care*, 26(1), 48-54.
19. Sun, K., Ren, M., Liu, D., Wang, C., Yang, C., & Yan, L. (2014). Alcohol consumption and risk of metabolic syndrome: a meta-analysis of prospective studies. *Clinical Nutrition*, 33(4), 596-602.
20. Wakabayashi, I. (2014). Frequency of heavy alcohol drinking and risk of metabolic syndrome in middle-aged men. *Alcoholism Clinical Experimental Research*, 38(6), 1689-1696.
21. Wei, M., Gibbons, L. W., Mitchell, T. L., Kampert, J. B., & Blair, S. N. (2000). Alcohol intake and incidence of type 2 diabetes in men. *Diabetes Care*, 23(1), 18-22.
22. Athyros, V. G., Liberopoulos, E. N., Mikhailidis, D. P., Papageorgiou, A. A., Ganotakis, E. S., Tziomalos, K., et al. (2007). Association of drinking pattern and alcohol beverage type with the prevalence of metabolic syndrome, diabetes, coronary heart disease, stroke, and peripheral arterial disease in a Mediterranean cohort. *Angiology*, 58(6), 689-697.
23. Chew, L. D., Nelson, K. M., Young, B. A., & Bradley, K. A. (2005). Association between alcohol consumption and diabetes preventive practices. *Family Medicine*, 37(8), 589-594.
24. Hodge, A. M., English, D. R., O'Dea, K., & Giles, G. G. (2006). Alcohol intake, consumption pattern and beverage type, and the risk of Type 2 diabetes. *Diabetic Medicine*, 23(6), 690-697.
25. Pietraszek, A., Gregersen, S., & Hermansen, K. (2010). Alcohol and type 2 diabetes. A review. *Nutrition, Metabolism and Cardiovascular Diseases*. doi:10.1016/j.numecd.2010.05.001.
26. Freiberg, M. S., Cabral, H. J., Heeren, T. C., Vasan, R. S., & Curtis Ellison, R. (2004). Alcohol consumption and the prevalence of the Metabolic Syndrome in the US.: a cross-sectional analysis of data from the Third National Health and Nutrition Examination Survey. *Diabetes Care*, 27(12), 2954-2959.
27. Gignoux, I., Gagnon, J., St-Pierre, A., Cantin, B., Dagenais, G. R., Meyer, F., et al. (2006). Moderate alcohol consumption is more cardioprotective in men with the metabolic syndrome. *The Journal of Nutrition*, 136(12), 3027-3032.
28. Rosell, M., De Faire, U., & Hellenius, M. L. (2003). Low prevalence of the metabolic syndrome in wine drinkers—is it the alcohol beverage or the lifestyle? *Eur J Clin Nutr*, 57(2), 227-234.
29. Alkerwi, A., Boutsen, M., Vaillant, M., Barre, J., Lair, M. L., Albert, A., et al. (2009). Alcohol consumption and the prevalence of metabolic syndrome: a meta-analysis of observational studies. *Atherosclerosis*, 204(2), 624-635.
30. Tortosa, A., Bes-Rastrollo, M., Sanchez-Villegas, A., Basterra-Gortari, F. J., Nunez-Cordoba, J. M., & Martinez-Gonzalez, M. A. (2007). Mediterranean diet inversely associated with the incidence of metabolic syndrome: the SUN prospective cohort. *Diabetes Care*, 30(11), 2957-2959.
31. Djousse, L., Arnett, D. K., Eckfeldt, J. H., Province, M. A., Singer, M. R., & Ellison, R. C. (2004). Alcohol consumption and metabolic syndrome: does the type of beverage matter? *Obesity Research*, 12(9), 1375-1385.

32. Baik, I., & Shin, C. (2008). Prospective study of alcohol consumption and metabolic syndrome. *American Journal of Clinical Nutrition*, 87(5), 1455-1463.
33. Wakabayashi, I. (2014). Frequency of heavy alcohol drinking and risk of metabolic syndrome in middle-aged men. *Alcoholism: Clinical and Experimental Research*, 38(6), 1689-1696.
34. Morrish, N. J., Wang, S. L., Stevens, L. K., Fuller, J. H., & Keen, H. (2001). Mortality and causes of death in the WHO Multinational Study of Vascular Disease in Diabetes. *Diabetologia*, 44 Suppl 2, S14-S21.
35. Gami, A. S., Witt, B. J., Howard, D. E., Erwin, P. J., Gami, L. A., Somers, V. K., et al. (2007). Metabolic syndrome and risk of incident cardiovascular events and death: a systematic review and meta-analysis of longitudinal studies. *Journal of the American College of Cardiology*, 49(4), 403-414.
36. Stern, M. P., Williams, K., Gonzalez-Villalpando, C., Hunt, K. J., & Haffner, S. M. (2004). Does the metabolic syndrome improve identification of individuals at risk of type 2 diabetes and/or cardiovascular disease? *Diabetes Care*, 27(11), 2676-2681.
37. Nanchahal, K., Ashton, W. D., & Wood, D. A. (2000). Alcohol consumption, metabolic cardiovascular risk factors and hypertension in women. *International Journal of Epidemiology*, 29(1), 57-64.
38. Avogaro, A., Watanabe, R. M., Dall'Arche, A., De Kreutzenberg, S. V., Tiengo, A., & Pacini, G. (2004). Acute alcohol consumption improves insulin action without affecting insulin secretion in type 2 diabetic subjects. *Diabetes Care*, 27(6), 1369-1374.
39. Bonnet, F., Disse, E., Laville, M., Mari, A., Hojlund, K., Anderwald, C. H., et al. (2012). Moderate alcohol consumption is associated with improved insulin sensitivity, reduced basal insulin secretion rate and lower fasting glucagon concentration in healthy women. *Diabetologia*, 55(12), 3228-3237.
40. Shai, I., Wainstein, J., Harman-Boehm, I., Raz, I., Fraser, D., Rudich, A., et al. (2007). Glycemic effects of moderate alcohol intake among patients with type 2 diabetes: a multicenter, randomized, clinical intervention trial. *Diabetes Care*, 30(12), 3011-3016.
41. Dixon, J. B., Dixon, M. E., & O'Brien, P. E. (2002). Alcohol consumption in the severely obese: relationship with the metabolic syndrome. *Obesity Research*, 10(4), 245-252.
42. Kastorini, C. M., Milionis, H. J., Esposito, K., Giugliano, D., Goudevenos, J. A., & Panagiotakos, D. B. (2011). The effect of Mediterranean diet on metabolic syndrome and its components: a meta-analysis of 50 studies and 534,906 individuals. *Journal of the American College of Cardiology*, 57(11), 1299-1313.
43. Volcik, K., Ballantyne, C. M., Pownall, H. J., Sharrett, A. R., & Boerwinkle, E. (2007). Interaction effects of high-density lipoprotein metabolism gene variation and alcohol consumption on coronary heart disease risk: The atherosclerosis risk in communities study. *Journal of Studies on Alcohol and Drugs*, 68(4), 485-492.
44. Brien, S. E., Ronksley, P. E., Turner, B. J., Mukamal, K. J., & Ghali, W. A. (2011). Effect of alcohol consumption on biological markers associated with risk of coronary heart disease: Systematic review and meta-analysis of interventional studies. *British Medical Journal*, 342, d636.
45. Beulens, J. W., van den Berg, R., Kok, F. J., Helander, A., Vermunt, S. H., & Hendriks, H. F. (2008). Moderate alcohol consumption and lipoprotein-associated phospholipase A2 activity. *Nutrition, Metabolism and Cardiovascular Diseases*, 18(8), 539-544.
46. Koppes, L. L., Dekker, J. M., Hendriks, H. F., Bouter, L. M., & Heine, R. J. (2006). Meta-analysis of the relationship between alcohol consumption and coronary heart disease and mortality in type 2 diabetic patients. *Diabetologia*, 49(4), 648-652.
47. Wakabayashi, I., Kobaba-Wakabayashi, R., & Masuda, H. (2002). Relation of drinking alcohol to atherosclerotic risk in type 2 diabetes. *Diabetes Care*, 25(7), 1223-1228.
48. Briasoulis, A., Agarwal, V., & Messerli, F. H. (2012). Alcohol consumption and the risk of hypertension in men and women: a systematic review and meta-analysis. *Journal of Clinical Hypertension (Greenwich)*, 14(11), 792-798.
49. Wakabayashi, I. (2011). Association between alcohol intake and metabolic syndrome in patients with hypertension. *Clinical and Experimental Hypertension*, 33(5), 299-303.
50. Bonaccio, M., Di Castelnuovo, A., Costanzo, S., Persichillo, M., De Curtis, A., Donati, M. B., et al. (2015). Adherence to the traditional Mediterranean diet and mortality in subjects with diabetes. Prospective results from the MOLI-SANI study. *European Journal of Preventive Cardiology*. doi:10.1177/2047487315569409.
51. Engler, P. A., Ramsey, S. E., & Smith, R. J. (2013). Alcohol use of diabetes patients: the need for assessment and intervention. *Acta Diabetologica*, 50(2), 93-99.
52. Djousse, L., Biggs, M. L., Mukamal, K. J., & Siscovick, D. S. (2007). Alcohol consumption and type 2 diabetes among older adults: The Cardiovascular Health Study. *Obesity*, 15(7), 1758-1765.
53. Carlsson, S., Hammar, N., Grill, V., & Kaprio, J. (2003). Alcohol consumption and the incidence of type 2 diabetes: a 20-year follow-up of the Finnish twin cohort study. *Diabetes Care*, 26(10), 2785-2790.
54. World Health Organization (WHO). (2013). *Global action plan for the prevention and control of noncommunicable diseases 2013-2020*. Geneva: World Health Organization.

Reviews

IARD Health & Policy Reviews cover the effects of alcohol consumption on health. They offer an overview of the relationship between drinking patterns and health outcomes, compile the key literature, and provide the reader with an extensive bibliography that refers to original research on each topic. The *Reviews* attempt to present the balance of the available evidence. They do not necessarily reflect the views of IARD or its sponsoring companies.

© International Alliance for Responsible Drinking (IARD) 2017. All Rights Reserved.

IARD is a not-for-profit organization dedicated to addressing the global public health issue of harmful drinking. Our mission is to contribute to the reduction of harmful drinking and promote responsible drinking worldwide. This is a problem that requires new insights, urgent action, and open dialogue. Central to IARD's work is our role as Secretariat of the Beer, Wine and Spirits Producers' Commitments to Reduce Harmful Drinking.



The International Alliance for
Responsible Drinking (IARD)
The Jefferson Building
1225 19th Street NW, Suite 500
Washington, DC 20036

IARD
INTERNATIONAL ALLIANCE FOR
RESPONSIBLE DRINKING

action on alcohol and global health

Phone: 1.202.986.1159
Fax: 1.202.986.2080
Email: info@iard.org
Web: www.iard.org